arXiv:hep-ph/9710523v1 29 Oct 1997

Exclusive Nonleptonic Decays of Heavy Baryons in a Relativistic Quark Model

V. E. Lyubovitskij[a,c], M. A. Ivanov,[a] J. G. Körner,[b] A. G. Rusetsky[a,d]

[a]JINR, Dubna, [b]Mainz University, [c]Tomsk University, [d]Tbilisi University

Abstract

Exclusive nonleptonic decays of bottom and charm baryons are studied within a relativistic quark model. We include factorizing as well as nonfactorizing contributions to the decay amplitudes.

Keywords: Nonleptonic decays; Heavy baryons; Quark models.
PACS: 12.39.Ki, 13.30.-a, 14.20.-c, 14.20.Lq, 14.20.Mr

In the near future one can expect large quantities of new data on exclusive charm and bottom baryon nonleptonic decays which calls for a comprehensive theoretical analysis of these decays. There exist a number of theoretical analysis of exclusive nonleptonic heavy baryon decays in the literature[1−6] including predictions for their angular decay distributions. The analysis of nonleptonic baryon decays is complicated by the necessity of having to include nonfactorizing contributions. In Fig.1 the Feynman diagrams which contribute to the amplitude of nonleptonic decays of heavy baryons are given: factorizing diagram (I), nonfactorizing diagrams (IIa, IIb, III). One thus has to go beyond the factorization approximation which had proved quite useful in the analysis of the exclusive nonleptonic decays of heavy mesons. There have been some theoretical attempts to analyse nonleptonic heavy baryon decays using factorizing contributions alone[5], the argument being that W-exchange contributions can be neglected in analogy to the power suppressed W-exchange contributions in the inclusive nonleptonic decays of heavy baryons. One might even be tempted to drop the nonfactorizing contributions on account of the fact that they are superficially proportional to $1/N_c$. However, since N_c-baryons contain N_c quarks an extra combinatorial factor proportional to N_c appears in the amplitudes which cancels the explicit diagrammatic $1/N_c$ factor[1,2]. There is now ample empirical evidences in the $c \to s$ sectors that nonfactorizing diagrams cannot be neglected. For example, in the charm sector the two observed decays $\Lambda_c^+ \to \Xi^0 K^+$ and $\Lambda_c^+ \to \Sigma\pi$ can only proceed via nonfactorizing diagrams. Their sizeable observed branching ratios may thus serve to obtain a measure of the size of the nonfactorizing contributions.

In this paper we use a relativistic quark model developed in Refs.[7] to calculate nonleptonic decays of heavy baryons. This model has been successfully applied to the description of the electromagnetic properties of nucleons[8] and has been extended to an analysis of the semileptonic decays of heavy baryons[9]. The detailed description of the model can be found in Refs.[7−9]. Since the evaluation of the nonleptonic decay amplitudes involves three-loop diagrams with nonlocal vertices (see Fig.1), we shall make some simplifying assumptions. These assumptions, on the one hand, have a clear physics motivation (we reproduce the analysis[1] of the spin structure of decay amplitudes in the spectator quark model[10,11]), and, on the other hand, allow one to evaluate both the factorizing and the nonfactorizing contributions to nonleptonic baryon decays. This can be achieved by assigning the projector $V_+ = (1+ \not{v})/2$ to each light quark field in the baryon-quark vertex, and by using the static approximation for u, d and s quark propagators in momentum space $S_q(k) = 1/m_q$ with m_q being a static quark mass. See, further details in Ref.[6]

Below we present our numerical results for decay rates and asymmetry parameters. Model parameters are determined from a fit to known branching ratios of nonleptonic decays $\Lambda_c^+ \to \Lambda^0\pi^+$, $\Lambda_c^+ \to \Sigma^0\pi^+$, $\Lambda_c^+ \to \Sigma^+\pi^0$, $\Lambda_c^+ \to p\bar{K}^0$ and $\Lambda_c^+ \to \Xi^0 K^+$ (see, Ref.[6]). The masses of hadrons are taken from Ref.[12]

In Tables 1 and 2 we give the results for the decay rates and asymmetry parameters for the heavy-to-light and $b \to c\bar{u}d$ nonleptonic decays, respectively.

Table 1. Branching ratios (in %) for heavy-light transitions.

Process	Körner, Krämer [1]	Xu, Kamal[3]	Cheng, Tseng[4]	Our	Experiment[12]
$\Lambda_c^+ \to \Lambda\pi^+$	0.76	1.67	0.91	0.79	0.79± 0.18
$\Lambda_c^+ \to \Sigma^0\pi^+$	0.33	0.35	0.74	0.88	0.88± 0.20
$\Lambda_c^+ \to \Sigma^+\pi^0$	0.33	0.35	0.74	0.88	0.88± 0.22
$\Lambda_c^+ \to \Sigma^+\eta$	0.16			0.11	0.48± 0.17
$\Lambda_c^+ \to \Sigma^+\eta'$	1.28			0.12	
$\Lambda_c^+ \to p\bar{K}^0$	2.16	1.24	1.30	2.06	2.2± 0.4
$\Lambda_c^+ \to \Xi^0 K^+$	0.27	0.10		0.31	0.34± 0.09
$\Xi_c^+ \to \Sigma^+\bar{K}^0$	5.11	0.35	0.67	3.08	
$\Xi_c^+ \to \Xi^0\pi^+$	2.80	2.66	3.12	4.40	1.2±0.5±0.3
$\Xi_c^0 \to \Lambda\bar{K}^0$	0.11	0.32	0.24	0.42	
$\Xi_c^0 \to \Sigma^0\bar{K}^0$	1.03	0.08	0.12	0.20	
$\Xi_c^0 \to \Sigma^+ K^-$	0.11	0.11		0.27	
$\Xi_c^0 \to \Xi^0\pi^0$	0.03	0.49	0.25	0.04	
$\Xi_c^0 \to \Xi^0\eta$	0.21			0.28	
$\Xi_c^0 \to \Xi^0\eta'$	0.74			0.31	
$\Xi_c^0 \to \Xi^-\pi^+$	0.91	1.52	1.10	1.22	
$\Omega_c^0 \to \Xi^0\bar{K}^0$	1.10		0.08	0.02	
$\Lambda_b^0 \to \Lambda\pi^0$				4.92×10^{-5}	
$\Lambda_b^0 \to pK^-$				2.11×10^{-4}	

Table 2. Branching ratios (in %) for heavy-heavy transitions.

Process	Γ (in 10^{10} s^{-1})	α	Process	Γ (in 10^{10} s^{-1})	α
$\Lambda_b^0 \to \Lambda_c^+\pi^-$	0.382	-0.99	$\Xi_b^0 \to \Xi_c'^0\pi^0$	0.014	0.94
$\Lambda_b^0 \to \Sigma_c^+\pi^-$	0.039	0.65	$\Xi_b^0 \to \Xi_c'^0\eta$	0.015	-0.98
$\Lambda_b^0 \to \Sigma_c^0\pi^0$	0.039	0.65	$\Xi_b^0 \to \Xi_c'^0\eta'$	0.021	0.97
$\Lambda_b^0 \to \Sigma_c^0\eta$	0.023	0.79	$\Xi_b^0 \to \Lambda_c^+ K^-$	0.010	-0.73
$\Lambda_b^0 \to \Sigma_c^0\eta'$	0.029	0.99	$\Xi_b^0 \to \Sigma_c^+ K^-$	0.030	-0.74
$\Lambda_b^0 \to \Xi_c^0 K^0$	0.021	-0.81	$\Xi_b^0 \to \Sigma_c^0\bar{K}^0$	0.021	0
$\Lambda_b^0 \to \Xi_c'^0 K^0$	0.032	0.98	$\Xi_b^0 \to \Omega_c^0 K^0$	0.023	0.65
$\Xi_b^0 \to \Xi_c^+\pi^-$	0.479	-1.00	$\Xi_b^- \to \Xi_c^0\pi^-$	0.645	-0.97
$\Xi_b^0 \to \Xi_c'^+\pi^-$	0.018	0.61	$\Xi_b^- \to \Xi_c'^0\pi^-$	0.007	-1.00
$\Xi_b^0 \to \Xi_c^0\pi^0$	0.002	-0.99	$\Xi_b^- \to \Sigma_c^0 K^-$	0.016	-0.98
$\Xi_b^0 \to \Xi_c^0\eta$	0.012	-0.86	$\Omega_b^- \to \Omega_c^0\pi^-$	0.352	0.60
$\Xi_b^0 \to \Xi_c^0\eta'$	0.003	0.71			

A clear pattern emerges. Nonfactorizing contributions plays a significant role in the heavy-to-light transitions. For heavy-to-heavy transitions the dominant rates are into channels with factorizing contributions. The rates which proceed only via nonfactorizing diagrams are small but not negligibly small.

In Tables 3 and 4 we analyse the contributions of nonfactorizing diagrams relative to those of the factorizing ones for the decays $\Lambda_c^+ \to \Lambda\pi^-$ and $\Lambda_b^0 \to \Lambda_c^+\pi^-$ which we predict to have the largest branching ratio. The total contribution of the nonfactorizing diagrams can be seen to be destructive. The sum of nonfactorizing contributions amount up to 60 % of the factorizing contribution in amplitude of heavy-to-light decay and up to 30 % in amplitude of $b \to c$ transition.

Table 3. Decay $\Lambda_c^+ \to \Lambda\pi^+$.
Contribution of nonfactorizing diagrams
(in % relative to the factorizing contribution)

Amplitude	Diagram			
	II_a	II_b	$II_a + II_b$	III
A	-29.8%	-18.5%	-48.3%	
B	-32.4%	-15.9%	-48.3%	-13.9%

Table 4. Decay $\Lambda_b^0 \to \Lambda_c^+\pi^-$.
Contribution of nonfactorizing diagrams
(in % relative to the factorizing contribution)

Amplitude	Diagram			
	II_a	II_b	$II_a + II_b$	III
A	-13.9%	-6.2%	-20.1%	
B	-14.3%	-5.8%	-20.1%	-8.5%



Fig. 1

Acknowledgements

M.A.I, V.E.L and A.G.R thank Mainz University for the hospitality where a part of this work was completed. This work was supported in part by the Heisenberg-Landau Program, by the Russian Fund of Basic Research (RFBR) under contract 96-02-17435-a, the State Committee of the Russian Federation for Education (project N 95-0-6.3-67, Grant Center at S.-Petersburg State University) and by the BMBF (Germany) under contract 06MZ566. J.G.K. acknowledges partial support by the BMBF (Germany) under contract 06MZ566.

References

1. J.G. Körner and M. Krämer, Z. Phys. **C55** (1992) 659.

2. J.G. Körner, in: Proc. VII Int. Conf. "Baryons '95" (Santa Fe, NM, Oct. 1995), eds. P.D. Barnes et al. (World Sci., Singapore, 1996) p. 221.

3. Q.P. Xu and A.N. Kamal, Phys. Rev. **D46** (1992) 270.

4. H.-Y. Cheng and B. Tseng, Phys. Rev. **D48** (1993) 4188.

5. H.-Y. Cheng, Phys. Rev. **D56** (1997) 2799.

6. M.A. Ivanov, J.G. Körner, V.E. Lyubovitskij and A.G. Rusetsky, Preprints MZ-TH/97-15, MZ-TH/97-21 (1997).

7. I.V. Anikin, M.A. Ivanov, N.B. Kulimanova and V.E. Lyubovitskij, Phys. Atom. Nucl. **57** (1994) 1082; Z. Phys. **C65** (1995) 681.
M.A. Ivanov and V.E. Lyubovitskij, Phys. Lett. **B408** (1997) 435.

8. M.A. Ivanov, M.P. Locher, V.E. Lyubovitskij, Few-Body Syst. **21** (1996) 131.

9. M.A. Ivanov, V.E. Lyubovitskij, J.G. Körner and P. Kroll, Phys. Rev. **D56** (1997) 348.

10. F. Hussain, J.G. Körner and G. Thompson, Ann. Phys. **206** (1991) 334.

11. F. Hussain, et al., Nucl. Phys. **B370** (1992) 259.

12. Review of Particle Physics, Phys. Rev., **D54** (1996) 1.